

Convert recurring decimals to fractions

Align the repeating tail, subtract, reduce to lowest terms and check reciprocals

• Recurring decimals and fraction form

01 Step 1: Identify the repeating block

- Distinguish a pure recurring decimal (0.232323...) from a mixed one (0.23333...).
- A reduced fraction p/q terminates only when q has no prime factors other than 2 and 5.
- Count non-repeating digits after the decimal point and digits in the repeating block.



02 Step 2: Multiply to align tails

- For $x = 0.232323\dots$, two digits repeat, so multiply by $10^2 = 100$ to get $100x = 23.232323\dots$
- Multiply y by 10 and 100 so both shifted forms, $10y = 2.333\dots$ and $100y = 23.333\dots$, share the same tail.
- For $y = 0.23333\dots$, one non-repeating digit precedes one repeating digit.



03 Step 3: Subtract and reduce

- Subtract aligned equations: $100x - x = 23$ gives $99x = 23$, so $x = 23/99$.
- Always reduce before summing terms: for $p/q = 23/99$ and $r/s = 7/30$, $p + q + r + s = 159$.
- For y : $100y - 10y = 21$ gives $90y = 21$, which reduces by gcd 3 to $y = 7/30$.



04 Step 4: Check proper and reciprocal forms

- For positive integers a, b : a/b is proper if $a < b$ and improper if $a \geq b$.
- Reciprocal b/a is proper only when $a > b$; thus $9/7$ has the proper reciprocal $7/9$.
- Inverting a non-zero fraction swaps numerator and denominator: reciprocal of a/b is b/a .

REMEMBER Shift decimals until identical repeating tails cancel on subtraction, then reduce by the GCD.